

# UNCLASSIFIED

AD NUMBER
AD435559
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; JAN 1964. Other requests shall be referred to Air Force Cambridge Research Laboratories, Hanscom AFB, MA.
AUTHORITY
AFCRL ltr dtd 3 Nov 1971

THIS PAGE IS UNCLASSIFIED

**UNCLASSIFIED**

**AD 435559**

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



**UNCLASSIFIED**

**NOTICE:** When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

AFCRL-64-16  
JANUARY 1964

435559

CATALOGED BY DDD

AS AD No. \_\_\_\_\_

AD 11-

Reprint



Research Report

"Reproduction in whole or in part  
is permitted for any purpose of  
the U. S. Government."

# **A Non-Destructive Radioanalytical Method for the Determination of Gallium in Substituted Yttrium Iron Garnets**

J. PAUL CALI

APR 14 1964

RECEIVED V. L. B.  
TICM. B

SOLID STATE SCIENCES LABORATORY PROJECT 5620

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES, OFFICE OF AEROSPACE RESEARCH, UNITED STATES AIR FORCE, L.G. HANSCOM FIELD, MASS.

NO. OTS

Requests for additional copies by Agencies of the Department of Defense, their contractors, and other government agencies should be directed to the:

Defense Documentation Center  
Cameron Station  
Alexandria, Virginia

Department of Defense contractors must be established for DDC services, or have their 'need-to-know' certified by the cognizant military agency of their project or contract.

(Reprinted from *Nature*, Vol. 199, No. 4898, pp. 1081-1082,  
September 14, 1963)

### A Non-destructive Radioanalytical Method for the Determination of Gallium in Substituted Yttrium Iron Garnets

THE substitution of diamagnetic ions with a valence of three for iron in garnets modifies both the saturation magnetization and the magnetic anisotropy of these materials. So far, there has been no correlation between the extent of the changes in these parameters and the amount of substituted ion. For these reasons and because the amount of garnet material was extremely limited, usually of the order of micrograms, a rapid, non-destructive method for the determination of gallium in yttrium iron garnet was undertaken. The empirical formula of this substituted garnet is  $(\text{Fe}_{1-x}\text{Ga}_x)\text{Y}_3\text{O}_{12}$ , where  $x$  is typically in the range of 0.05-0.25.




Fig. 1. Gamma spectra of three samples as indicated. Intensities are relative. Spectra taken 4 h after irradiation. A, Gallium yttrium iron garnet; B, gallium standard; C, yttrium iron garnet

Attempts to measure the gallium content by X-ray fluorescence methods failed because of the anomalous results obtained between single crystals and powders of the same composition. A non-destructive radioanalytical scheme based on neutron activation and  $\gamma$ -ray pulse-height analysis was developed in this laboratory.

Gallium exists as two stable isotopes, gallium-69 and gallium-71. Thermal neutron irradiation of the former produces the  $\beta$ -emitter (99+ per cent) gallium-70, but because  $\gamma$ -spectrometry was to be used for measurement of the induced activity, it was not of interest in this work. Under the same irradiation conditions, gallium-72 is also formed. Its principal  $\gamma$ -lines are found in good yield at 0.63 and 0.83 MeV; its half-life was found to be 14.2 h, in good agreement with reported values<sup>1</sup>.

The spherically shaped, gallium-doped garnet samples weighing from 200 to 300 micrograms were placed in an irradiation container together with a suitable amount of pure gallium metal and a garnet (YIG) containing no gallium. Samples and standards were then irradiated in the pneumatic facility of the Massachusetts Institute of Technology Reactor for 1 min at a thermal flux of  $8 \times 10^{11}$  neutrons  $\text{cm}^{-2} \text{sec}^{-1}$ .

The induced  $\gamma$ -activity was analysed for both energy and decay characteristics using a 512-channel analyser coupled to a 2-in. well-type sodium iodide (TI) crystal detector.

Fig. 1 shows some typical results. The upper curve, A, is the  $\gamma$ -spectrum of the gallium-doped garnet. The middle curve, B, is that due to the pure gallium metal, and the lowest curve, C, is that of the pure gallium-free yttrium iron garnet. The principal peaks in the two upper curves are found to be at 0.63 and 0.83 MeV in agreement with the literature<sup>2</sup>  $\gamma$ -energy values for gallium-72.

Quantitative results are obtained by measurement of the areas under the samples and standard 0.83-MeV peaks after applying the usual background, decay, aliquoting and geometry corrections. An unusual correction, possible because of the versatility of the equipment used herein, was the subtraction of the activity under the aforementioned peak due to the pure yttrium iron garnet. This correction ranged from 2 to 4 per cent. Some results obtained are shown in Table 1.

Table 1. TYPICAL RESULTS OF THE RADIOACTIVATION ANALYSIS OF GALLIUM-SUBSTITUTED YTTRIUM IRON GARNETS

	C.p.m./ $\mu\text{g}$ in the 0.83 $\gamma$ peak*	Per cent gallium by weight
Sample A	61.5	12.2 $\pm$ 0.6
Sample B	50.4	10.0 $\pm$ 0.5
Sample C	47.9	9.5 $\pm$ 0.5
Standard	506	

\* All corrections applied; time of determination was 4 h after irradiation.

Statistical, weighing, and other known errors amounted individually to less than 1 per cent, and the standard error of the method, given as the standard deviation, is calculated to be not more than 5 per cent.

A search of the literature<sup>3</sup> indicates that no single interfering element or combination of elements could give a spectrogram with  $\gamma$ -energies and decay characteristics similar to that found.

Finally, it should be pointed out that because of the short irradiation time, residual activity, after a few days decay, is minimal and the garnets may be recovered for additional investigations.

J. PAUL CALI

JOSEPH R. WEINER

Air Force Cambridge Research Laboratories,  
Office of Aerospace Research,  
L. G. Hanscom Field, Bedford, Mass.

JOHN J. O'CONNOR

Tracerlab, a Division of Laboratory for Electronics, Inc.,  
Waltham, Mass.

<sup>1</sup> Koch, R. C., *Activation Analysis Handbook* (Academic Press, New York, 1960).

<sup>2</sup> Crouthamel, C. E., *Applied Gamma-Ray Spectrometry* (Pergamon, London, 1960).

<sup>3</sup> Strominger, D., Hollander, J. M., and Seaborg, G. T., *Rev. Modern Phys.*, **30**, 585 (1958).